arXiv:1202.1998v2 [stat.ME] 28 Mar 2012

Hierarchical Kendall copulas: Properties and inference

Eike Christian Brechmann*

This version: March 27, 2012

First version: November 2, 2011

Abstract

While there is substantial need for dependence models in high dimensions, most existing models strongly suffer from the curse of dimensionality and barely balance parsimony and flexibility. In this paper, the new class of hierarchical Kendall copulas is proposed which tackles these problems. Constructed with flexible copulas specified for groups of variables in different hierarchical levels, hierarchical Kendall copulas are able to model complex dependence patterns without severe restrictions. The paper explicitly discusses inference techniques for hierarchical Kendall copulas, in particular, simulation, estimation and model selection. A substantive application to German stock returns finally shows that hierarchical Kendall copulas perform very well, out-of- as well as in-sample.

Keywords: multivariate copula, hierarchical copula, Kendall distribution function

1 Introduction

Dependence modeling using copulas has made significant progress in the last years. Many of the standard, and also of the newly proposed, models however suffer from the curse of dimensionality, which makes it virtually impossible to use them for very large data sets as required, e.g., in financial or spatial applications. One common approach to overcome the curse of dimensionality is grouping data, e.g., by industry sectors or nationality. Such copula models include the grouped Student-t copula by Daul et al. (2003) and hierarchical Archimedean copulas, which were initially proposed by Joe (1997). In particular, hierarchical structures such as the latter are very appealing and received considerable attention lately (see, e.g., Hofert (2010)).

A major issue of any copula model is to find a good balance between parsimony and flexibility. While elliptical copulas such as the common Gaussian and Student-t require an enormous number of parameters for specifying the correlation matrix (the number of parameter grows quadratically with the dimension), Archimedean and also hierarchical Archimedean copulas are much more parsimonious, since the number of parameters is at most linear in the dimension. However, such restrictions may be severe, since hierarchical Archimedean copulas are at the same time limited to the class of Archimedean copulas as building blocks.

The purpose of this paper is to introduce the new class of hierarchical Kendall copulas, which is a flexible but yet parsimonious copula model available for applications also in higher dimensions. It is built up by copulas for groups of variables in different hierarchical levels.

---

*Center for Mathematical Sciences, Technische Universität München, Boltzmannstr. 3, D-85747 Garching, Germany. E-mail: brechmann@ma.tum.de. Phone: +49 89 289-17425. Fax: +49 89 289-17435.

In particular—and in contrast to hierarchical Archimedean copulas—, it does not require any restrictions with respect to copula choice or the parameters and therefore is able to model complex patterns of dependence such as asymmetry and tail dependence between large numbers of variables.

The name "hierarchical Kendall copula" is chosen to stress the central role of the Kendall distribution function in the model formulation. The Kendall distribution function is the distribution function of the multivariate probability integral transform using the copula. It is used to aggregate the (dependence) information of a group of variables. It was first studied in Genest and Rivest (1993) in the bivariate case and in more detail in Barbe et al. (1996). Other accounts on it can be found, e.g., in Imlahi et al. (1999), Chakak and Imlahi (2001), Genest and Rivest (2001) and Nelsen et al. (2003) as well as in the copula goodness-of-fit literature (see, e.g., Wang and Wells (2000)).

It has been shown by Genest et al. (1995) that the only copula which gives a valid multivariate distribution for non-overlapping multivariate marginals is the independence copula. Marco and Ruiz-Rivas (1992) state conditions how a distribution function with specified multivariate marginals can be constructed; the easiest case being that margins are max-infinitely divisible, which includes distributions based on Archimedean copulas. Hierarchical Kendall copulas circumvent such issues through aggregation facilitated by the Kendall distribution functions. This implies that in general the obtained multivariate distribution is not a copula.

The model, which I call hierarchical Kendall copula, has previously been mentioned by Anjos and Kolev (2005), who however do not further develop the model in terms of statistical properties and inference. The work presented here is completely independent of theirs and develops in detail how hierarchical Kendall copulas can be used for statistical inference; simulation, estimation and model selection are discussed explicitly. Finally, a substantial 30-dimensional application to German stock returns is presented, showing the good out-of- as well as in-sample performance of hierarchical Kendall copulas. Hierarchical Kendall copulas therefore answer the need for parsimonious and flexible models in high dimensions and provide means to overcome the curse of dimensionality in dependence modeling.

The remainder of the paper is organized as follows. The new model is introduced and discussed in Section 2. Section 3 then treats statistical inference technique for hierarchical Kendall copulas, where an extensive simulation study is performed to validate the estimation methods. In Section 4 the financial application is presented in detail, while Section 5 concludes.

2 Hierarchical Kendall copulas

A central part of the definition of hierarchical Kendall copulas, which will be given below, is the notion of the Kendall distribution function which therefore is treated first. Subsequently, hierarchical Kendall copulas are defined and their properties are discussed, in particular in contrast to hierarchical Archimedean copulas.

2.1 Kendall distribution functions

Kendall distribution functions were first studied in two dimensions by Genest and Rivest (1993) and studied in more generality by Barbe et al. (1996). They are defined as follows.

Definition 2.1 (Kendall distribution function). *Let C be a d-dimensional copula. For* $\boldsymbol{U} :=$

$(U_1, ..., U_d) \sim C$ *the Kendall distribution function is defined as*

$$K^{(d)}(t) := P(C(\boldsymbol{U}) \leq t), \quad t \in (0,1). \tag{2.1}$$

In other words, the Kendall distribution function is the univariate distribution function of the random variable $Z := C(\boldsymbol{U})$, the multivariate probability integral transform of $\boldsymbol{U}$. It has been shown by Genest and Rivest (1993) that bivariate Archimedean copulas are uniquely characterized by their Kendall distribution functions. Genest et al. (2011) recently extended this result to the trivariate case and strongly conjecture that this holds in general.

For my purposes, a specific interpretation of Kendall distribution functions will be particularly useful. First, I define the univariate function

$$C_{u_1,...,u_{d-1}}(u_d) := C(u_1, ..., u_{d-1}, u_d)$$

and denote the corresponding inverse as $C^{-1}_{u_1,...,u_{d-1}}$ which is the *quantile function* of a copula as studied in Imlahi et al. (1999) and Chakak and Ezzerg (2000), since

$$C(u_1, ..., u_{d-1}, C^{-1}_{u_1,...,u_{d-1}}(z)) = z$$

for $z \in (0,1)$. This function will be useful for describing the *level set* of a copula

$$L(z) = \{\boldsymbol{u} \in [0,1]^d : C(\boldsymbol{u}) = z\}, \quad z \in (0,1), \tag{2.2}$$

which is closely related to the Kendall distribution function, since

$$P(L(z)) = P(C(\boldsymbol{U}) = z) =: k^{(d)}(z),$$

if the Kendall distribution function $K^{(d)}$ is differentiable with density $k^{(d)}$.

For ease of notation I further define for $r \leq d-1$

$$C_{u_1,...,u_r}(u_d) := C_{u_1,...,u_r,1,...,1}(u_d),$$

and for $z \in (0,1)$

$$C^{-1}_{\emptyset}(z) := z.$$

Example 2.2 (Quantile function of an exchangeable Archimedean copula)**.** *For a d-dimensional exchangeable Archimedean copula (see McNeil and Nešlehová (2009)) with generator*[1] φ*,*

$$C(u_1, ..., u_d) = \varphi^{-1}\left(\sum\nolimits_{1 \leq i \leq d} \varphi(u_i)\right),$$

the quantile function can by determined as

$$C^{-1}_{u_1,...,u_{d-1}}(z) = \varphi^{-1}\left(\varphi(z) - \sum\nolimits_{1 \leq i < d} \varphi(u_i)\right).$$

Using the above notation, the following theorem provides a way to compute Kendall distribution functions of arbitrary copulas.

---

[1] The function φ generates a d-dimensional exchangeable Archimedean copula if and only if its inverse φ^{-1} is d-monotone on $[0,\infty)$, that is, φ^{-1} is differentiable up to the order $d-2$ on $[0,\infty)$, it holds that $(-1)^k(\varphi^{-1})^{(k)}(x) \geq 0$ for $k = 0, 1, ..., d-2$ and for any $x \in [0,\infty)$, and $(-1)^{d-2}(\varphi^{-1})^{(d-2)}$ is non-increasing and convex on $[0,\infty)$ (see McNeil and Nešlehová (2009)).

Family	Generator $\varphi(t)$	Kendall dist. function $K(t)$
Independence	$-\log(t)$	$t\,(1-\log(t))$
Clayton	$t^{-\theta}-1, \quad \theta > 0$	$t\left(1+\frac{1-t^{\theta}}{\theta}\right)$
Gumbel	$(-\log(t))^{\theta}, \quad \theta \geq 1$	$t\left(1-\frac{\log(t)}{\theta}\right)$
Frank	$-\log\left(\frac{1-e^{-\theta t}}{1-e^{-\theta}}\right), \quad \theta \in \mathbb{R}\setminus\{0\}$	$t+\frac{e^{\theta t}-1}{\theta}\log\left(\frac{1-e^{-\theta}}{1-e^{-\theta t}}\right)$
Joe	$-\log(1-(1-t)^{\theta}), \quad \theta > 1$	$t-\frac{\log(1-(1-t)^{\theta})(1-(1-t)^{\theta})}{\theta(1-t)^{\theta-1}}$
Countermonotonic	$1-t$	1
Comonotonic	-	t

Table 1: Generator and Kendall distribution functions of bivariate Archimedean copulas as well as the Kendall distribution functions of the bivariate Fréchet-Hoeffding lower and upper bounds.

Theorem 2.3 (Kendall distribution function of a d-dimensional copula)**.** *Let* $\boldsymbol{U} := (U_1, ..., U_d) \sim C$*, where* C *is absolutely continuous with density* c*. If* $d = 2$*, the Kendall distribution function* $K^{(2)}$ *is given by (Chakak and Ezzerg 2000)*

$$K^{(2)}(t) = t + \int_t^1 \int_0^{C_{u_1}^{-1}(t)} c(u_1, u_2)du_2du_1 = t + \int_t^1 \frac{\partial}{\partial u_1} C(u_1, u_2)\Big|_{u_2 = C_{u_1}^{-1}(t)} du_1.$$

Following Imlahi et al. (1999) the Kendall distribution function of a d*-dimensional copula can then be obtained recursively through*

$$K^{(d)}(t) = K^{(d-1)}(t) + \int_t^1 \int_{C_{u_1}^{-1}(t)}^1 ... \int_{C_{u_1,...,u_{d-2}}^{-1}(t)}^1 \int_0^{C_{u_1,...,u_{d-1}}^{-1}(t)} c(u_1, ..., u_d)\ du_d...du_1,$$

where $K^{(d)}$ *denotes the Kendall distribution function of the* d*-dimensional copula and* $K^{(d-1)}$ *that of the* $(d-1)$*-dimensional margin of the first* $(d-1)$ *variables.*

For general copulas it is not possible to easily determine the Kendall distribution function in closed form. A convenient exception are however (exchangeable) Archimedean copulas.

Theorem 2.4 (Kendall distribution function of an Archimedean copula)**.** *Let* $\boldsymbol{U} := (U_1, ..., U_d) \sim C$*, where* C *is an exchangeable Archimedean copula. Then the Kendall distribution function is given by (Barbe et al. 1996)*

$$K^{(d)}(t) = t + \sum_{i=1}^{d-1} (-1)^i \frac{(\varphi(t))^i}{i!} (\varphi^{-1})^{(i)}(\varphi(t)).$$

In particular, if $d = 2$*,*

$$K^{(2)}(t) = t - \frac{\varphi(t)}{\varphi'(t)} =: t - \lambda(t),$$

with the λ*-function first studied in Genest and Rivest (1993).*

Table 1 shows generator and Kendall distribution functions of common bivariate Archimedean copulas (see Joe (1997) and Nelsen (2006)) as well as of the Fréchet-Hoeffding lower and upper bounds. In case of the independence copula, the Kendall distribution function for general dimensions also has a simple closed-form expression.



Figure 1: Kendall distribution functions of the independence copula (2.3) (left panel) and the Gumbel copula with parameter $\theta = 2$ (right panel) for different dimensions d.

Example 2.5 (Kendall distribution function of the d-dimensional independence copula)**.** *The Kendall distribution function of the d-dimensional independence copula is conveniently given as*

$$K^{(d)}(t) = t + t \sum_{i=1}^{d-1} \frac{(\log(1/t))^i}{i!}. \tag{2.3}$$

Example 2.5 shows that the Kendall distribution function of the independence copula is, for fixed t, increasing with the dimension d. Since Kendall distribution functions are obviously bounded from above by 1—which in two dimensions is the Kendall distribution function of the Fréchet-Hoeffding lower bound—, the Kendall distribution function of the independence copula converges to 1 as $d \to \infty$. This is illustrated in the left panel of Figure 1.

For comparison the right panel of Figure 1 shows the Kendall distribution function of the Gumbel copula with parameter $\theta = 2$ (medium positive dependence). While here the Kendall distribution function also converges to 1 as $d \to \infty$, this convergence is much slower. The practical implications of this property on my model will be discussed below.

In the following, the superscript of the Kendall distribution function, which indicates the dimension of the associated random vector, will be omitted for reasons of readability.

2.2 Model formulation and properties

The definition of the new dependence model class of hierarchical Kendall copulas is now stated. Although the model has previously been formulated by Anjos and Kolev (2005), it has—to the best of my knowledge—not yet been treated in detail or used for statistical inference. I choose the name *hierarchical Kendall copula* to stress the central role the Kendall distribution function plays in the model formulation.

Definition 2.6 (Hierarchical Kendall copula)**.** *Let $U_1, ..., U_n \sim U(0, 1)$. Further let $C_0, C_1, ..., C_d$ be copulas of dimension d for C_0 and $1 \leq n_i \leq d$ for C_i, $i \in \{1, ..., d\}$, where $n = \sum_{i=1}^{d} n_i$, respectively. Moreover let $K_1, ..., K_d$ denote the Kendall distribution functions corresponding to $C_1, ..., C_d$, respectively, and define $m_i = \sum_{j=1}^{i} n_j$ for $i = 0, ..., d$. Then the hierarchical Kendall copula is defined as follows:*

(i) $(U_{m_{i-1}+1}, ..., U_{m_i}) \sim C_i \ \forall i \in \{1, ..., d\}$,

(ii) $(V_1, ..., V_d) \sim C_0$, *where* $V_i := K_i(C_i(U_{m_{i-1}+1}, ..., U_{m_i})) \ \forall i \in \{1, ..., d\}$.


$\boldsymbol{U}_1 := (U_1, ..., U_{m_1}) \sim C_1$
$\boldsymbol{U}_2 := (U_{m_1+1}, ..., U_{m_2}) \sim C_2$
$\cdots$
$\boldsymbol{U}_d := (U_{m_{d-1}+1}, ..., U_{m_d}) \sim C_d$
$C_1(\boldsymbol{U}_1) \sim K_1$
$C_2(\boldsymbol{U}_2) \sim K_2$
$\cdots$
$C_d(\boldsymbol{U}_d) \sim K_d$
$V_1 := K_1(C_1(\boldsymbol{U}_1)) \sim U(0,1)$
$V_2 := K_2(C_2(\boldsymbol{U}_2)) \sim U(0,1)$
$\cdots$
$V_d := K_d(C_d(\boldsymbol{U}_d)) \sim U(0,1)$
$(V_1, ..., V_d) \sim C_0$

Figure 2: Illustration of the hierarchical Kendall copula model of Definition 2.6.

Definition 2.6 is illustrated in Figure 2. Note that C_0 is in general *not* a copula of $(U_1, ..., U_n)$ but of $(V_1, ..., V_d)$, which are uniform random variables, since $C_i(U_{m_{i-1}+1}, ..., U_{m_i}) \sim K_i \ \forall i \in \{1, ..., d\}$. Typically C_0 is denoted as *nesting copula*, while $(U_{m_{i-1}+1}, ..., U_{m_i})$, $i \in \{1, ..., d\}$, is called *cluster*. The nesting copula C_0 and the cluster copulas $C_1, ..., C_d$ can be chosen independently. They can be arbitrary copulas such as common Archimedean or elliptical copulas or from any other class of copulas. From now on it is assumed that they are *absolutely continuous* and possess densities c_0 and $c_1, ..., c_d$, respectively.

The Kendall distribution functions $K_1, ..., K_d$ are used to summarize the information contained in the clusters (transformation of n_i-variate to univariate random variable). While one may also think of other transformations, I believe that Kendall distribution functions are particularly useful for this purpose. First, since the Kendall distribution function is monotone, C_0 essentially models the comovement of the contour levels (2.2). This can be seen as a proxy for the strength of dependence in the clusters, since the dimensionality of the single clusters is “normalized” through the Kendall distribution functions. Second, other variables such as components of an elliptical distribution do not contain such specific information which summarizes the information in a single variable. And third, a focus on tails, as it is often desired, is empirically hardly feasible, since tail behavior is very hard to quantify appropriately. Thus, the transformation using the Kendall distribution function reasonably summarizes the (dependence) information of a multivariate random vector. For the particular purpose of risk aggregation an alternative hierarchical dependence model for sums of random variables was recently proposed by Arbenz et al. (2011).

The two-level construction given in Definition 2.6 can easily be defined for an arbitrary number of levels.

Definition 2.7 (Hierarchical Kendall copula with k levels)**.** *Let* $U_1, ..., U_n \sim U(0,1)$ *and* $d_1 \geq d_2 \geq ... \geq d_{k-1}$*. Further let* C_0 *be a* d_{k-1}*-dimensional copula and* $C_i^{(j)}$, $j = 1, ..., k-1$ *(index running over the* k *levels),* $i = 1, ..., d_j$ *(index running over the* d_j *clusters of each level* j*), be copulas of dimension* $1 \leq n_i^{(j)} \leq d_j$*, where* $n = \sum_{i=1}^{d_1} n_i^{(1)}$ *and* $d_{j-1} = \sum_{i=1}^{d_j} n_i^{(j)}$ *for* $j = 2, ..., k-1$*. Further let* $K_i^{(j)}$ *denote the Kendall distribution function corresponding to* $C_i^{(j)}$, $j = 1, ..., k-1, i = 1, ..., d_j$*, and define* $m_i^{(j)} = \sum_{j=1}^{i} n_j^{(j)}$ *for* $i = 0, ..., d$*. Then the hierarchical Kendall copula with* k *levels is defined as follows:*

(i) $(U_{m_{i-1}^{(1)}+1}, ..., U_{m_i^{(1)}}) \sim C_i^{(1)} \ \forall i \in \{1, ..., d_1\}$,


$\boldsymbol{U}_1 := (U_1, ..., U_{m_1^{(1)}}) \sim C_1^{(1)}$
$\cdots$
$\boldsymbol{U}_\ell := (U_{m_{\ell-1}^{(1)}+1}, ..., U_{m_\ell^{(1)}}) \sim C_\ell^{(1)}$
$\cdots$
$V_1^{(1)} := K_1(C_1(\boldsymbol{U}_1)) \sim U(0,1)$
$\cdots$
$V_\ell^{(1)} := K_\ell(C_\ell(\boldsymbol{U}_\ell)) \sim U(0,1)$
$\cdots$
$\boldsymbol{V}_1 := \left(V_1^{(1)}, ..., V_{m_1^{(2)}}^{(1)}\right) \sim C_1^{(2)}$
$\cdots$
$\boldsymbol{V}_{d_2} := \left(V_{m_{d_2-1}^{(2)}+1}^{(1)}, ..., V_{m_{d_2}^{(2)}}^{(1)}\right) \sim C_{d_2}^{(2)}$
$V_1^{(2)} := K_1(C_1(\boldsymbol{V}_1)) \sim U(0,1)$
$\cdots$
$V_{d_2}^{(2)} := K_{d_2}(C_{d_2}(\boldsymbol{V}_{d_2})) \sim U(0,1)$
$(V_1^{(2)}, ..., V_d^{(2)}) \sim C_0$

Figure 3: Illustration of a three level hierarchical Kendall copula model (see Definition 2.7). Here $m_1^{(2)} = \ell$.

(ii) $V_i^{(1)} := K_i^{(1)}(C_i^{(1)}(U_{m_{i-1}^{(1)}+1}, ..., U_{m_i^{(1)}}))\ \forall i \in \{1, ..., d_1\}$,

(iii) for $j = 2, ..., k-1$*:*

  (a) $(V^{(j-1)}_{m_{i-1}^{(j)}+1}, ..., V^{(j-1)}_{m_i^{(j)}}) \sim C_i^{(j)}\ \forall i \in \{1, ..., d_j\}$,

  (b) $V_i^{(j)} := K_i^{(j)}(C_i^{(j)}(V^{(j-1)}_{m_{i-1}^{(j)}+1}, ..., V^{(j-1)}_{m_i^{(j)}})\ \forall i \in \{1, ..., d_j\}$,

(iv) $(V_1^{(k-1)}, ..., V_{d_{k-1}}^{(k-1)}) \sim C_0$.

An example of a three level hierarchical Kendall copula is shown in Figure 3. For simplicity and illustrative reasons I restrict my exposition here to the case of $k = 2$ hierarchical levels. All derivations and methods described in the following can easily be generalized to the general k level case.

The joint density function of a hierarchical Kendall copula is now derived under certain independence assumptions.

Theorem 2.8 (Joint density of a hierarchical Kendall copula)**.** *Let* $\boldsymbol{U} = (U_1, ..., U_n)$ *be distributed according to a hierarchical Kendall copula. Further let* $\boldsymbol{U}_i = (U_{m_{i-1}+1}, ..., U_{m_i})$ *and* $\boldsymbol{u}_i = (u_{m_{i-1}+1}, ..., u_{m_i})$*,* $i = 1, ...d$*. Similarly* $\boldsymbol{u} = (u_1, ..., u_n)$*. If*

$$\boldsymbol{U}_i \perp \boldsymbol{U}_j | (V_1, ..., V_d) \quad \textit{and} \quad \boldsymbol{U}_i \perp V_j | V_i \quad \forall i \neq j,\ i, j \in \{1, ..., d\}, \tag{2.4}$$

then the joint density function $c_{\mathcal{K}}$ *of a hierarchical Kendall copula is given as follows:*

$$c_{\mathcal{K}}(\boldsymbol{u}) = c_0(K_1(C_1(\boldsymbol{u}_1)), ..., K_d(C_d(\boldsymbol{u}_d))) \prod_{i=1}^{d} c_i(\boldsymbol{u}_i). \tag{2.5}$$

Proof: By adding only redundant information, one gets

$$\begin{aligned} c_{\mathcal{K}}(\boldsymbol{u}) &= P(\boldsymbol{U} = \boldsymbol{u}) = P(\boldsymbol{U} = \boldsymbol{u}, V_1 = K_1(C_1(\boldsymbol{u}_1)), ..., V_d = K_d(C_d(\boldsymbol{u}_d))) \\ &= P(\boldsymbol{U} = \boldsymbol{u} | V_j = K_j(C_j(\boldsymbol{u}_j)), j = 1, ..., d) P(V_j = K_j(C_j(\boldsymbol{u}_j)), j = 1, ..., d), \end{aligned} \tag{2.6}$$

where $V_j := K_j(C_j(\boldsymbol{U}_j)),\ j = 1, ..., d$. Following Definition 2.6 the second term of Equation (2.6) is then given by

$$P(V_j = K_j(C_j(\boldsymbol{u}_j)), j = 1, ..., d) = c_0(K_1(C_1(\boldsymbol{u}_1)), ..., K_d(C_d(\boldsymbol{u}_1))). \tag{2.7}$$

On the other hand, the first term of Equation (2.6) simplifies to

$$\begin{aligned} P(\boldsymbol{U} = \boldsymbol{u} | V_j = K_j(C_j(\boldsymbol{u}_j)), j = 1, ..., d) &= \prod_{i=1}^{d} P(\boldsymbol{U}_i = \boldsymbol{u}_i | V_j = K_j(C_j(\boldsymbol{u}_1)), j = 1, ..., d) \\ &= \prod_{i=1}^{d} P(\boldsymbol{U}_i = \boldsymbol{u}_i | V_i = K_i(C_i(\boldsymbol{u}_i))), \end{aligned}$$

because of condition (2.4) which in particular implies that $\boldsymbol{U}_i|(V_1, ..., V_d) = \boldsymbol{U}_i|V_i$.

Further,

$$P(\boldsymbol{U}_i = \boldsymbol{u}_i | V_i = K_i(C_i(\boldsymbol{u}_i))) = P(\boldsymbol{U}_i = \boldsymbol{u}_i) = c_i(\boldsymbol{u}_i) \quad \forall i = 1, ..., d, \tag{2.8}$$

since the conditioning is redundant information.

The proof concludes by plugging Equations (2.7) and (2.8) into Equation (2.6). □

I do not regard condition (2.4), which perfectly fits into the setting of Definition 2.6, as rather restrictive. The intuition behind the two assumptions is that, given the information of the nesting variables $V_1, ..., V_d$, the clusters are independent of each other and also of other nesting variables, since the dependence among the clusters is explained through the "representatives" $V_1, ..., V_d$. In numerical experiments this was confirmed empirically.

Theorem 2.8 now allows to state the following corollary which summarizes the marginal properties of hierarchical Kendall copulas.

Corollary 2.9 (Margins of a hierarchical Kendall copula)**.** *The same notation as in Theorem 2.8 is used.*

(i) Bivariate margins: Let $k, \ell \in \{1, ..., n\},\ k \neq \ell$. W.l.o.g. $k < \ell$.

  (a) If U_k and U_ℓ are in the same cluster i, their marginal distribution function $C_{k\ell}$ is

$$C_{k\ell}(u_k, u_\ell) := C_i(1, ..., 1, u_k, 1, ..., 1, u_\ell, 1, ..., 1).$$

  (b) If U_k and U_ℓ are in different clusters and condition (2.4) holds, their marginal distribution function $C_{k\ell}$ is

$$\begin{aligned} C_{k\ell}(u_k, u_\ell) := \int_0^{u_k} \int_0^{u_\ell} \int_{[0,1]^{n-2}} & c_{\mathcal{K}}(u_1, ..., u_{k-1}, w_1, u_{k+1}, ..., u_{\ell-1}, w_2, u_{\ell+1}, ..., u_n) \\ & du_1 ... du_{k-1} du_{k+1} ... du_{\ell-1} du_{\ell+1} ... du_n dw_2 dw_1. \end{aligned} \tag{2.9}$$

(ii) Multivariate margins: The marginal distribution of the cluster $\boldsymbol{U}_i$ is C_i.

Proof: Claims *(i)(a)* and *(ii)* directly follow from Definition 2.6. With regard to *(i)(b)*, Theorem 2.8 yields the bivariate marginal density $c_{k\ell}$ of U_k and U_ℓ as

$$c_{k\ell}(u_k, u_\ell) = \int_{[0,1]^{n-2}} c_{\mathcal{K}}(\boldsymbol{u})\ du_1 ... du_{k-1} du_{k+1} ... du_{\ell-1} du_{\ell+1} ... du_n, \tag{2.10}$$

from which Equation (2.9) follows through integration. □

More general multivariate margins involving variables from different clusters can be derived as in Equation (2.9).

Furthermore bivariate marginal distributions where the variables are in different clusters can be regarded as some kind of continuous mixture of the nesting copula C_0. By plugging (2.5) into (2.10), the density of $C_{k\ell}$ as defined above is given by

$$c_{k\ell}(u_k, u_\ell) = \int_{[0,1]^{n-2}} c_0(K_1(C_1(\boldsymbol{u}_1)), ..., K_d(C_d(\boldsymbol{u}_d))) \prod_{i=1}^{d} c_i(\boldsymbol{u}_i)$$
$$du_1 ... du_{k-1} du_{k+1} ... du_{\ell-1} du_{\ell+1} ... du_n,$$

where the mixing density weights are given by $\prod_{i=1}^{d} c_i(\boldsymbol{u}_i)$.

2.3 Hierarchical Kendall copulas vs. hierarchical Archimedean copulas

The popular class of hierarchical Archimedean copulas also allows for a nested modeling of clusters of variables. In contrast to hierarchical Kendall copulas, hierarchical Archimedean copulas are however limited to Archimedean copulas as building blocks, while hierarchical Kendall copulas can be built up by any possible copula. Furthermore, hierarchical Kendall copulas do not have any restrictions regarding the dependence, while hierarchical Archimedean copulas require stronger intra-cluster dependence, which results in parameter restrictions if generators are the same.

As will be discussed below, hierarchical Kendall copulas also benefit from many of the convenient properties of Archimedean copulas. It will be shown that closed-form sampling of hierarchical Kendall copulas is feasible, when cluster copulas are (exchangeable) Archimedean. Hierarchical Kendall copulas with Archimedean cluster copulas are further particularly easy to estimate, since Kendall distribution functions are known in closed form for Archimedean copulas. For that reason they also provide a closed-form joint density function (under condition (2.4)), which is numerically tractable even in higher dimensions.

In contrast to hierarchical Archimedean copulas the nesting copula C_0 is however not closed under addition and removal of cluster components U_ℓ. This is because the Kendall distribution function is not independent with respect to the dimension (cp. Theorem 2.3). That is, if a random variable U_{n+1} is added to cluster $j \in \{1, ..., d\}$, the transformation K_j and thus V_j change, even if C_j is exchangeable Archimedean. Similarly, if a random variable is removed from a cluster.

3 Inference of hierarchical Kendall copulas

In the following, statistical inference techniques for hierarchical Kendall copulas are discussed. First simulation is treated, then estimation and model selection.

3.1 Simulation

The following general sampling procedure describes how to obtain one single sample from a given hierarchical Kendall copula.

Algorithm 3.1 (Simulation of hierarchical Kendall copulas)**.** *Let* $C_0, C_1, ..., C_d$ *be known. Further let* K_i^{-1} *denote the inverse of the Kendall distribution function* K_i *for* $i = 1, ..., d$ *and assume that it is also known.*

(i) Obtain a sample $(v_1, ..., v_d)$ *from* C_0*.*

(ii) Set $z_i := K_i^{-1}(v_i)\ \forall i \in \{1, ..., d\}$*.*

(iii) Obtain a sample $(u_{m_{i-1}+1}, ..., u_{m_i})$ *from* $(U_{m_{i-1}+1}, ..., U_{m_i})|C_i(U_{m_{i-1}+1}, ..., U_{m_i}) \equiv z_i$ *for* $i = 1, ..., d$*.*

(iv) Return $\boldsymbol{u} := (u_1, ..., u_n)$*.*

Given that one can simulate from the copula C_0, sampling from hierarchical Kendall copulas thus amounts to the more general question of sampling from a distribution $\boldsymbol{U}|C(\boldsymbol{U}) \equiv z$, where C is the copula of a marginally uniform random vector $\boldsymbol{U} := (U_1, ..., U_d)$ and $z \in (0, 1)$. In other words, one wants to sample from a multivariate distribution given a specific level set (2.2) at level z as illustrated in the left panel of Figure 4. This problem is discussed in the following three sections, where the first section solves the issue using (approximate) rejection sampling, while the second provides expressions for conditional distribution functions, which are shown to be available in closed form for exchangeable Archimedean copulas. The third section provides an alternative solution for Archimedean copulas, which is based on recent results from the literature.

Before moving on, I however like to note that in contrast to the assumptions in Algorithm 3.1 the Kendall and the copula distribution functions may not be known in general. This problem is treated explicitly in Section 3.2, where estimation of hierarchical Kendall copulas is discussed. When Kendall distribution functions are known, inverses are typically obtained by numerical inversion, since closed-form expressions are often not available.

3.1.1 Rejection sampling

To obtain an approximate sample from $\boldsymbol{U}|C(\boldsymbol{U}) \equiv z$, rejection sampling can be used as in Imlahi et al. (1999): Instead of sampling from $\boldsymbol{U}|C(\boldsymbol{U}) \equiv z$, one chooses a small number $\varepsilon > 0$ and sample from $\boldsymbol{U}|(z - \varepsilon \leq C(\boldsymbol{U}) \leq z + \varepsilon)$. This is illustrated in the middle panel of Figure 4.

Algorithm 3.2 (Rejection sampling)**.** *Let* $z \in (0, 1)$ *and* $\varepsilon > 0$*.*

(i) Obtain a sample $\boldsymbol{u} := (u_1, ..., u_d)$ *from* C*.*

(ii) If $|C(\boldsymbol{u}) - z| < \varepsilon$*, return* $\boldsymbol{u}$*. Otherwise go back to step (i).*

The right panel of Figure 4 shows an exemplary sample from a Clayton copula with parameter $\theta = 2$ at contour level $z = 0.2$. For illustrative reasons ε is chosen as 0.01.

Algorithm 3.2 can efficiently be combined with Algorithm 3.1 in order to simultaneously obtain N samples from a hierarchical Kendall copula.



Figure 4: Left panel: observations from a bivariate Clayton copula with parameter $\theta = 2$ and level sets at levels $z = 0.1, ..., 0.9$ (solid lines). Middle panel: level set at level $z = 0.2$ of the bivariate Clayton copula distribution function with parameter $\theta = 2$ (solid line). The dashed lines illustrate the rejection sampling area $[z - 0.01, z + 0.01]$. Right panel: scatter plot of corresponding samples generated through rejection sampling.

Algorithm 3.3 (Simulation of hierarchical Kendall copulas using rejection sampling)**.** *Let* C_0*,* $C_1, ..., C_d$ *and* $K_1^{-1}, ..., K_d^{-1}$ *be known. Further let* $\varepsilon > 0$ *be given.*

- *(i) Obtain* N *samples* $(v_{j1}, ..., v_{jd})$*,* $j = 1, ..., N$*, from* C_0*.*
- *(ii) For* $i = 1, ..., d$*:*
  - *(a) Set* $z_{ji} := K_i^{-1}(v_{ji})$ $\forall j \in \{1, ..., N\}$ *and define* $D := \{1, ..., N\}$*.*
  - *(b) Obtain a sample* $\tilde{\boldsymbol{u}} := (\tilde{u}_{m_{i-1}+1}, ..., \tilde{u}_{m_i})$ *from* C*.*
  - *(c) If for any* $j' \in \{1, ..., N\}$*,* $|C(\tilde{\boldsymbol{u}}) - z_{j'i}| < \varepsilon$*, set* $(u_{j',m_{i-1}+1}, ..., u_{j',m_i}) := \tilde{\boldsymbol{u}}$ *and* $D := D \setminus \{j'\}$*.*[2] *Go back to step (b) until* $D = \emptyset$*.*
- *(iii) Return samples* $(u_{j1}, ..., u_{jn})$*,* $j = 1, ..., N,$*.*

The computational efficiency obviously depends on the choice of ε. The smaller ε is, the longer Algorithm 3.3 takes, but, at the same time, the more accurate the results are. Due to rounding errors in simulation and calculation of the copula distribution function, it is in fact reasonable to allow for an error ε.

3.1.2 Conditional inverse method

A common method to generate samples from a multivariate distribution is the conditional inverse method (see Devroye (1986)). For this one needs to determine the iterative conditional distribution functions of $\boldsymbol{U}|C(\boldsymbol{U}) \equiv z$, that is, of $U_1|C(\boldsymbol{U}) \equiv z$, $U_2|(U_1 = u_1, C(\boldsymbol{U}) \equiv z)$, ..., $U_d|(U_1 = u_1, ..., U_{d-1} = u_{d-1}, C(\boldsymbol{U}) \equiv z)$.[3] The corresponding conditional distribution functions are denoted by $F_{j|1,...,j-1}(\cdot|u_1, ..., u_{j-1}, z)$ and densities by $f_{j|1,...,j-1}(\cdot|u_1, ..., u_{j-1}, z)$ for $j = 1, ..., d$, respectively. Then one obtains a sample from $\boldsymbol{U}|C(\boldsymbol{U}) \equiv z$ using the following algorithm.

---

[2] If there is more than one sample j with $|C(\tilde{\boldsymbol{u}}) - z_{ji}| < \varepsilon$, choose j' such that $|C(\tilde{\boldsymbol{u}}) - z_{j'i}| < |C(\tilde{\boldsymbol{u}}) - z_{ji}|$ $\forall j$.

[3] The distribution of $U_d|(U_1 = u_1, ..., U_{d-1} = u_{d-1}, C(\boldsymbol{U}) \equiv z)$ actually does not need to be determined because the value of U_d is uniquely given through the conditioning variables.

Algorithm 3.4 (Conditional inverse method)**.** *Let* $z \in (0,1)$*.*

- *(i) Obtain* $(d-1)$ *uniform samples* $v_1, ..., v_{d-1}$*.*
- *(ii) For* $j = 1, ..., d-1$*:* $u_j := F^{-1}_{j|1,...,j-1}(v_j|u_1, ..., u_{j-1}, z)$*.*
- *(iii) Set* $u_d := C^{-1}_{u_1,...,u_{d-1}}(z)$*.*
- *(iv) Return sample* $(u_1, ..., u_d)$*.*

The problem however is to determine the conditional distribution functions which are generally not given in closed form.

Theorem 3.5 (Conditional distributions)**.** *Let* $\boldsymbol{U} \sim C$*, then* $\forall j = 1, ..., d$

$$F_{j|1,...,j-1}(u|u_1, ..., u_{j-1}, z) = \frac{\int_{C^{-1}_{u_1,...,u_{j-1}}(z)}^{u} g(u_j)\ du_j}{\int_{C^{-1}_{u_1,...,u_{j-1}}(z)}^{1} g(u_j)\ du_j}, \quad u \in (C^{-1}_{u_1,...,u_{j-1}}(z), 1), \tag{3.1}$$

where

$$g(u_j) = \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} c(u_1, ..., u_{d-1}, C^{-1}_{u_1,...,u_{d-1}}(z)) \frac{\partial}{\partial z} C^{-1}_{u_1,...,u_{d-1}}(z)\ du_{d-1}...du_{j+1}. \tag{3.2}$$

The proof can be found in Appendix A.

Evidently the conditional distribution functions given in (3.1) in general do not allow for explicit expressions. In particular, if C^{-1} is not available in closed form such as for the Gaussian copula, the expression in (3.1) hardly simplifies.

In the case of exchangeable Archimedean copulas the conditional distribution functions can however be obtained in closed form.

Lemma 3.6 (Conditional distributions of exchangeable Archimedean copulas)**.** *Let* $\boldsymbol{U} \sim C$*, where* C *is a* d*-dimensional exchangeable Archimedean copula, then* $\forall j = 1, ..., d$

$$F_{j|1,...,j-1}(u_j|u_1, ..., u_{j-1}, z) = \left(1 - \frac{\varphi(u_j)}{\varphi(z) - \sum_{1 \leq i < j} \varphi(u_i)}\right)^{d-j}, \quad u \in (C^{-1}_{u_1,...,u_{j-1}}(z), 1). \tag{3.3}$$

The proof can again be found in Appendix A.

Lemma 3.6 then allows to restate Algorithm 3.4 for exchangeable Archimedean copulas, for which the conditional distribution functions can easily be inverted in closed form.

Algorithm 3.7 (Conditional inverse method for exchangeable Archimedean copulas)**.** *Let* C *be an exchangeable Archimedean copula with generator* φ *and* $z \in (0,1)$*.*

- *(i) Obtain* $(d-1)$ *uniform samples* $v_1, ..., v_{d-1}$*.*
- *(ii) For* $j = 1, ..., d-1$*:* $u_j := \varphi^{-1}((1 - v_j^{1/(d-j)})(\varphi(z) - \sum_{1 \leq i < j} \varphi(u_i)))$*.*
- *(iii) Set* $u_d := \varphi^{-1}(\varphi(z) - \sum_{1 \leq i < d} \varphi(u_i))$*.*
- *(iv) Return sample* $(u_1, ..., u_d)$*.*



Figure 5: Left panel: scatter plot of a sample of a bivariate Clayton copula with parameter $\theta = 2$ at $z = 0.2$. Middle and right panel: 3D scatter plot and pairs plot of a sample of a trivariate Clayton copula with parameter $\theta = 2$ at $z = 0.2$.

Step *(iii)* could also be performed in step *(ii)*. It is however not necessary to use an additional uniform sample v_d. This is why steps *(ii)* and *(iii)* are shown separately.

For illustration Figure 5 shows scatter plots of samples from bivariate and trivariate Clayton copulas with parameter $\theta = 2$. Notice in particular the difference to the right panel of Figure 4.

As a side note, one observes that Algorithm 3.7 can in particular be used to sample from a given exchangeable Archimedean copula. This is described in the following algorithm of which an equivalent version has previously been stated in Wu et al. (2007). The idea is to first simulate the contour level and then sample conditional on that level set, since

$$P(\boldsymbol{U} = \boldsymbol{u}) = P(C(\boldsymbol{U}) = z)P(\boldsymbol{U} = \boldsymbol{u} | C(\boldsymbol{U}) = z).$$

Algorithm 3.8 (Sampling from exchangeable Archimedean copulas)**.** *Let C be an exchangeable Archimedean copula with generator φ and Kendall distribution function K (see Theorem 2.4), whose inverse K^{-1} is assumed to exist.*

(i) Obtain d uniform samples $v_1, ..., v_d$.

(ii) Set $z := K^{-1}(v_d)$.

(iii) For $j = 1, ..., d$: $u_j := \varphi^{-1}((1 - v_j^{1/(d-j)})(\varphi(z) - \sum_{1 \leq i < j} \varphi(u_i)))$.

(iv) Return sample $(u_1, ..., u_d)$.

3.1.3 Projected distribution

Clearly the condition $C(\boldsymbol{U}) \equiv z$ on the distribution of $\boldsymbol{U}$ means that one is in fact investigating a $(d-1)$-dimensional distribution, namely the distribution of $\boldsymbol{U}$ projected onto the level set $C(\boldsymbol{U}) \equiv z$ which is a $(d-1)$-dimensional manifold in $[0,1]^d$. However this distribution is not easily tractable in general.

Again in the case of exchangeable Archimedean copulas some convenient results are obtainable. McNeil and Nešlehová (2009) show that if $\boldsymbol{U} \sim C$, where C is an exchangeable Archimedean copula with generator φ, then

$$(\varphi(U_1), ..., \varphi(U_d)) \overset{d}{=} R\boldsymbol{S},$$

where $\boldsymbol{S} = (S_1, ..., S_d)$ is uniformly distributed on the unit simplex $\{\boldsymbol{x} \geq 0 : \sum_{j=1}^{d} x_k = 1\}$, a $(d-1)$-dimensional manifold in $[0,1]^d$, and the *radial part*

$$R = \sum_{j=1}^{d} \varphi(U_j)$$

has distribution F_R which can be determined through the *Williamson transform* (see McNeil and Nešlehová (2009)).

Recall that $C(\boldsymbol{u}) = \varphi^{-1}(\sum_{j=1}^{d} \varphi(u_j))$ holds for exchangeable Archimedean copulas. Fixing the contour level $C(\boldsymbol{U}) \equiv z$ is thus equivalent to setting $R \equiv \varphi(z)$, so that

$$[\boldsymbol{U} | C(\boldsymbol{U}) \equiv z] \overset{d}{=} \left(\varphi^{-1}(S_1 \varphi(z)), ..., \varphi^{-1}(S_d \varphi(z))\right).$$

This result can then be used to provide another sampling algorithm for $\boldsymbol{U} | C(\boldsymbol{U}) \equiv z$, which can be shown to be equivalent to Algorithm 3.7, when using explicit expressions for the sample $(s_1, ..., s_d)$ from $\boldsymbol{S}$ in terms of uniform random variables (see Hering (2011, Lemma 3.1.8)).

Algorithm 3.9 (Projected distribution sampling for exchangeable Archimedean copulas)**.** *Let C be an exchangeable Archimedean copula with generator φ and $z \in (0,1)$.*

(i) Obtain a sample $(s_1, ..., s_d)$ from $\boldsymbol{S}$.

(ii) For $j = 1, ..., d$: $u_j := \varphi^{-1}(s_j \varphi(z))$.

(iii) Return sample $(u_1, ..., u_d)$.

When also sampling the contour level to obtain observations from the Archimedean copula, this is again the approach proposed by Wu et al. (2007) which has been restated by Hering (2011) in the setting of the work by McNeil and Nešlehová (2009) as I use it here.

As stated above, it is an open problem to determine this projected distribution for general copulas. Analogous sampling methods to Algorithm 3.9 could be used then.

3.2 Estimation and model selection

Estimation of model parameters in hierarchical Kendall copulas can either be performed stepwise or jointly when the joint density expression is available. Both estimation methods are discussed here and evaluated and compared in an extensive simulation study. Finally some remarks regarding model selection are made.

3.2.1 Two-step estimation

The hierarchical construction given in Definition 2.6 directly leads to a two-step estimation procedure of hierarchical Kendall copulas.

Algorithm 3.10 (Two-step estimation of hierarchical Kendall copulas)**.** *Let $(\boldsymbol{u_j})_{j=1,...,N}$ be a sample of a hierarchical Kendall copula as defined in Definition 2.6 ($\boldsymbol{u_j} = (u_{j1}, ..., u_{jn})$ for $j = 1, ..., N$). Further let $\boldsymbol{\theta}_0, \boldsymbol{\theta}_1, ..., \boldsymbol{\theta}_d$ be the parameters of the copulas $C_0, C_1, ..., C_d$, respectively. One then obtains corresponding estimates $\hat{\boldsymbol{\theta}}_i$, $i = 0, ..., d$, as follows.*

(i) For $i = 1, ..., d$ estimate $\boldsymbol{\theta}_i$ based on $(u_{j,m_{i-1}+1}, ..., u_{j,m_i})_{j=1,...,N}$.

(ii) Estimate $\boldsymbol{\theta}_0$ *based on* $v_{ji} := K_i(C_i(u_{j,m_{i-1}+1}, ..., u_{j,m_i}|\hat{\boldsymbol{\theta}}_i)|\hat{\boldsymbol{\theta}}_i),\ i = 0, ..., d, j = 1, ..., N.$

The Kendall distribution functions $K_i,\ i = 1, ..., d$, depend on $\hat{\boldsymbol{\theta}}_i$ through C_i as indicated by the notation.

As briefly noted before, inference of hierarchical Kendall copulas poses two main challenges: In order to move up the hierarchy, the copula distribution functions C_i and the Kendall distribution function K_i have to be known for all $i = 1, ..., d$. While this is the case for (exchangeable) Archimedean copulas, this is not true in general. In particular, for the popular class of elliptical copulas neither the copula nor the Kendall distribution functions are known in closed form.

These two issues can be tackled as follows: If the copula distribution function C_i is unknown, one can substitute it by the corresponding empirical version

$$\hat{C}_i(u_{m_{i-1}+1}, ..., u_{m_i}) = \frac{1}{N}\sum_{j=1}^{N} \mathbf{1}_{\{u_{j,m_{i-1}+1} \leq u_{m_{i-1}+1}, ..., u_{j,m_i} \leq u_{m_i}\}},$$

since possibly high-dimensional integration of the density expression is not feasible in general. The accuracy of the empirical copula can be improved by obtaining a large sample of the copula under consideration. Other non-parametric estimators for the copula distribution functions may be used but are potentially computationally more demanding.

Having computed transformed data

$$z_{ji} := C_i(u_{j,m_{i-1}+1}, ..., u_{j,m_i}|\hat{\boldsymbol{\theta}}_i),\ i = 1, ..., d, j = 1, ..., N,$$

one possibility to get an approximation of the Kendall distribution function K_i is simply to take the empirical distribution function of $(z_{ji})_{j=1,...,N}$, i.e.,

$$\hat{K}_i(z) = \frac{1}{N}\sum_{j=1}^{N} \mathbf{1}_{\{z_{ji} \leq z\}}. \tag{3.4}$$

Alternatively I propose to use a *kernel* estimator

$$\hat{K}_i(z) = \frac{1}{N}\sum_{j=1}^{N} G\left(\frac{z - z_{ji}}{h}\right), \tag{3.5}$$

where G is a kernel distribution function and h an appropriately chosen bandwidth (see, e.g., Li and Racine (2006)). If few observations are available, this approach yields a smoother function. In particular, if n_i is large, K_i may be close to 1 (see Figure 1) so that a smoother estimate may be helpful to capture the properties of the distribution function evaluated close to 0. Other possible approaches may be *splines* (see, e.g., Xue and Wang (2010)).

If one is only interested in the nesting copula, one can simply use the non-parametric methods described above for the clusters and then only estimate the parameters of the nesting copula. Moreover this two-step estimation procedure directly generalizes to a k-step estimation approach for k level hierarchical Kendall copulas as defined in Definition 2.7.

3.2.2 Joint estimation

Given the joint density function (2.5) of a hierarchical Kendall copula (assuming that condition (2.4) holds), parameters can also be estimated jointly using the joint log likelihood expression

$$\ell_{\mathcal{K}}(\boldsymbol{\theta}_0, \boldsymbol{\theta}_1, ..., \boldsymbol{\theta}_d) = \sum_{j=1}^{N} \log c_{\mathcal{K}}(\boldsymbol{u}_j|\boldsymbol{\theta}_0, \boldsymbol{\theta}_1, ..., \boldsymbol{\theta}_d). \tag{3.6}$$

As starting values for a joint maximum likelihood estimation, stepwise estimates as described above can be used.

For this to be practically feasible the copula and Kendall distribution functions of the cluster copulas have to been known. If this is not the case, empirical versions can be used to approximate the joint log likelihood expression. When the joint log likelihood expression is however available common model selection techniques such as AIC and BIC can be used to check the model fit.

3.2.3 Simulation study

In order to validate the estimation procedures introduced above, I performed a large scale Monte Carlo study. For this I simulated from a four-dimensional hierarchical Kendall copula (two bivariate clusters) and then estimated the parameters according to the following methods:

- Sequential (two-step) estimation with known Kendall distribution functions.
- Sequential estimation with empirical Kendall distribution functions (3.4).
- Sequential estimation with kernel estimator (3.5) for the Kendall distribution functions (Epanechnikov kernel).
- MLE with known starting values (true parameters).
- MLE with sequentially estimated starting values.

The cluster copulas C_1 and C_2 were chosen as Clayton, Gumbel or Frank; the nesting copula C_0 as Gaussian, Clayton, Gumbel or Frank. Parameters were determined according to Kendall's τ values of 0.4 and 0.7. Sample sizes were 250, 500 and 1000 and the number of repetitions was 100.

MSEs of the estimated nesting copula parameter θ_0 (transformed to Kendall's τ values) according to the above five procedures are shown in Figure 6 for the case of Clayton and Gumbel copulas; the other cases are not shown here due to lack of space and since it turns out that results are essentially independent of the chosen cluster copulas.

Similarly, the choice of parameters of the cluster copulas does not have a great influence on the results, while larger nesting parameters mean more accurate results. The results are however not independent of the nesting copula. For the Frank nesting copula, there is hardly any difference between the five estimation procedures, but overall, MLE and sequential estimation with known Kendall distribution functions are best. When the Kendall distribution functions are unknown, the standard empirical estimator is superior to the kernel estimator of the Kendall distribution functions in the case of a Clayton nesting copula and vice versa if the nesting copula is a Gaussian or a Gumbel. Finally, while MSEs are generally rather small, there also is a clear positive effect of an increasing sample size.

To summarize, especially when the Kendall distribution functions are known, estimation of hierarchical Kendall copulas is very accurate. When they are not known, no clear recommendation can be made whether to use a standard empirical or a kernel estimator.

3.2.4 Model selection

In practical applications the clusters $(U_{m_{i-1}+1}, ..., U_{m_i}), i \in \{1, ..., d\}$, have to be identified. In cases where they are not given from the data (e.g., industry sectors in financial data), common clustering techniques (see, e.g., Hastie et al. (2009)) can be used after having specified a



Figure 6: MSEs of estimated parameters in terms of Kendall's τ values of the five estimation procedures for different sample sizes n and nesting copulas C_0. Cluster copula 1: Clayton. Cluster copula 2: Gumbel. Denotation of x-axis: (τ_0,τ_1,τ_2), where $L := 0.4$ and $H := 0.7$. y-axes are chosen such that MSEs in each column are comparable.

Sectors	Fin.	Chem.	Healtc.	Auto.	Ind.	Retail	IT	Util.	Transp.
Mean pairw. Kendall's τ	0.41	0.33	0.21	0.39	0.38	0.26	0.28	0.56	0.29
Est. deg. of freedom	9.01	10.97	24.41	12.96	8.85	10.67	9.08	4.75	7.37

Table 2: Mean pairwise empirical Kendall's τ and estimated degrees of freedom of a Student-t copula for each cluster.

measure of closeness among variables. If a multi-level hierarchical Kendall copula is considered, hierarchical clustering methods may be particularly helpful.

Recalling the discussion at the end of Section 2.1 the size of clusters has to be carefully chosen because Kendall distributions may become almost degenerate at 0 for very large clusters. In most practical situations this is however not an issue, since already under medium positive dependence the convergence to the constant function at 1 is very slow (see the right panel of Figure 1).

4 Application

Finance is a major field, where copulas are used for dependence modeling (see, e.g., Cherubini et al. (2004)). Often financial data exhibits some kind of clustering structure such as industry sectors and national stock markets. For such data, hierarchical Kendall copulas are very suitable. To investigate the usefulness of this newly proposed class of dependence models and to illustrate the presented inference techniques, the most important German stock market index DAX are analyzed.

The DAX is composed of 30 major German stocks. For these I identified ten industry sectors: financials (Allianz, Commerzbank, Deutsche Bank, Deutsche Börse, Munich Re), chemicals (BASF, Bayer, K+S, Linde), healthcare (Fresenius, Fresenius Medical Care, Merck), automobile (BMW, Daimler, Volkswagen), industrials (MAN, Siemens, ThyssenKrupp), retail and consumer goods (Adidas, Beiersdorf, Henkel, Metro), IT and communications (Deutsche Telekom, Infineon, SAP), utilities (E.ON, RWE), transportation and logistics (Deutsche Post, Lufthansa), and building materials (HeidelbergCement).

For all 30 stocks more than six years of log returns (January 2005 to July 2011) are considered, where the time series are separated into a training set of 1158 observations and a testing set of 500 observations for out-of-sample validation of my models. As it is common for copula modeling in finance, I preliminarily fit time series models to the marginal time series and then work with standardized residuals which are transformed to marginally uniform data by the probability integral transform (inference functions from margins (IFM) method by Joe (1997)). In particular, marginal GARCH(1,1)-models with Student-t innovations are chosen, which have been validated with appropriate tests.

Table 2 shows the mean pairwise empirical Kendall's τ and the estimated degrees of freedom of a multivariate Student-t copula for each cluster. Evidently, within-sector dependence is variable, since some clusters are more homogeneous than others. Also strong tail dependence, as indicated by small degrees of freedom, cannot be found in all clusters.

I then fitted different hierarchical Kendall copulas to the training data set. Results (log likelihood, AIC, BIC) are reported in Table 3. As building block copulas I considered Clayton, Gumbel and Frank (Archimedean) as well as Gaussian and Student-t (elliptical) copulas to

Model	# Par.	2-Step Loglik.	Joint MLE Loglik.	Increase	AIC	BIC
(Clayton, Student-t)	55	6593.28	6677.73	1.2%	-13245.46	-12967.47
(Gumbel, Student-t)	55	6980.12	6992.29	0.2%	-13874.57	-13596.58
(Frank, Student-t)	55	7178.55	7190.29	0.2%	-14270.59	-13992.59
(Gaussian, Student-t)	82	7587.05	-	-	-15010.09	**-14595.63**
(Student-t, Student-t)	91	7854.31	-	-	**-15526.61**	**-15066.66**
(Clayton, Clayton)	10	5452.34	5471.09	0.3%	-10922.17	-10871.63
(Gumbel, Gumbel)	10	5860.74	5862.93	0.0%	-11705.85	-11655.31
Multivariate Gaussian	435	8422.09	-	-	**-15974.19**	-13775.50
Multivariate Student-t	436	8861.91	-	-	**-16851.82**	**-14648.08**

Table 3: Estimation results. Notation of models: (cluster copulas, nesting copula). Correlation matrices of multivariate Gaussian and Student-t copulas are estimated using inversion of pairwise empirical Kendall's τ's, while the degrees of freedom parameter is estimated via MLE given that correlation matrix. AIC and BIC values are based on the joint MLE if available (best three values are indicated in bold).

account for different dependedence structure. The specified models are shown in the first column of Table 3. Note that no results of models with Gaussian nesting copulas are displayed. This is because those models always performed inferior to respective models with Student-t copula.

For comparison I also fitted multivariate Gaussian and Student-t copulas, for which however no joint MLE was possible due to the large number of parameters. Also no joint MLE for hierarchical Kendall copulas with Gaussian and Student-t cluster copulas was performed due to uncertainty in the Kendall distribution function. Since two-step estimation provides good estimates (see column "Increase" in Table 3), this is however not a severe limitation.

A 30-dimensional hierarchical Archimedean copula could not be fitted for comparison, since its density expression is hardly tractable and numerically very involved (see Savu and Trede (2010)). A stepwise approach by fitting clusters first and then the nesting Archimedean copula fails due to the dependence restrictions of hierarchical Archimedean copulas (see Section 2.3): while there is moderate dependence within some clusters (cp. Table 2), there still is considerable dependence among clusters, which cannot be modeled using a hierarchical Archimedean copula.

The hierarchical Kendall copulas therefore benefit from not having such dependence restrictions. Moreover, ten-dimensional Student-t nesting copulas appear more reasonable than exchangeable Archimedean nesting copulas (with only one parameter). With respect to cluster copulas, symmetric copulas (Frank, Gaussian, Student-t) are superior to asymmetric ones (Clayton, Gumbel). The larger flexibility when using Gaussian and Student-t copulas is reflected in smaller information criteria, although more parameters are required. In comparison to standard multivariate Gaussian and Student-t copulas, hierarchical Kendall copulas perform quite well, in particular when taking into account the enormous number of parameters of these models.

4.1 Value-at-Risk forecasting

In finance, interest is however not so much in a good in-sample fit but rather in out-of-sample validation. A typical exercise for this is Value-at-Risk (VaR) forecasting. If the distribution of returns is continuous, the $(1-\alpha)$-VaR is the α-quantile of the distribution. For risk management this value needs to be predicted on a daily basis, which I do for the training set of 500 days using moving windows of length 1158.

Forecasts are typically evaluated in terms of *exceedances*, that is, the event that the predicted

Model	Level	# Exceed.	UC	IND1	IND2	CC1	CC2	CC3	CC4
Independence copula	99%	103	*0.00*	*0.04*	*0.00*	*0.00*	*0.00*	*0.00*	*0.00*
	95%	135	*0.00*	0.15	*0.03*	*0.00*	*0.00*	*0.00*	*0.00*
	90%	157	*0.00*	0.17	*0.03*	*0.00*	*0.00*	*0.00*	*0.00*
Gaussian copula	99%	5	1.00	0.75	0.79	0.95	0.35	0.93	1.00
	95%	26	0.84	0.09	0.74	0.23	0.35	0.98	1.00
	90%	52	0.77	0.46	0.87	0.73	*0.03*	0.29	0.73
Student-t copula	99%	4	0.64	0.80	0.21	0.87	0.93	0.75	0.95
	95%	24	0.84	0.12	0.73	0.29	0.39	0.89	1.00
	90%	53	0.66	0.53	0.85	0.74	0.05	0.31	0.73
Kendall (Gaussian)	99%	5	1.00	0.75	0.82	0.95	0.35	0.93	1.00
	95%	27	0.69	0.08	0.72	0.20	0.31	0.97	1.00
	90%	56	0.38	0.46	0.84	0.51	*0.03*	0.23	0.65
Kendall (Frank)	99%	2	0.13	0.90	0.46	0.31	0.35	0.43	0.82
	95%	31	0.23	0.04	0.76	0.06	0.21	0.55	0.92
	90%	57	0.31	0.52	0.81	0.48	*0.03*	0.22	0.64
Kendall (Student-t)	99%	5	1.00	0.75	0.80	0.95	0.35	0.93	1.00
	95%	28	0.55	0.07	0.74	0.16	0.32	0.90	1.00
	90%	57	0.31	0.52	0.78	0.48	*0.03*	0.22	0.64

Table 4: P-values of VaR backtests for hypotheses of independence and (un)conditional coverage. Hierarchical Kendall copulas use a Student-t nesting copula and cluster copulas as indicated in brackets.

VaR is exceeded by the observed return. For 500 forecasts on average $500 \times \alpha$ exceedances are expected. Whether the number of exceedances (*unconditional coverage*) and their occurrences (should be *independent*; both properties: *conditional coverage*) are appropriate can be evaluated using a range of tests (*backtests*) that have been proposed in the literature: the proportion of failures test of unconditional coverage by Kupiec (1995) (UC), the Markov test of independence by Christoffersen (1998) (IND1), the joint test of conditional coverage by Christoffersen (1998) (CC1), the mixed Kupiec test of conditional coverage by Haas (2001) (CC2), the Weibull test of independence by Christoffersen and Pelletier (2004) (IND2), and the duration-based GMM test of conditional coverage by Candelon et al. (2011) (CC3 and CC4 with orders 2 and 5, respectively).

Here, the Value-at-Risk of an equally weighted portfolio of the 30 DAX stocks is forecasted. Backtesting results of the following six different models can be found in Table 4:

- independence copula (for comparison),
- Gaussian and Student-t copulas,
- hierarchical Kendall copulas with Gaussian, Frank and Student-t cluster copulas and with Student-t nesting copula.

In summary, none of the hypotheses of independence and (un)conditional coverage can consistently be rejected for any of the VaR levels and for any of the models—except for the multivariate independence copula as expected. The weak lack of conditional coverage at the 90% level as detected by the mixed Kupiec test of Haas (2001) is not supported by the other tests.

This shows that hierarchical Kendall copulas are as good as the common Gaussian and Student-t copulas when it comes to out-of-sample validation. In contrast to these models, the hierarchical Kendall copulas are however much more parsimonious and have straightforward interpretation in terms of sectorial dependence. In particular, the hierarchical Kendall copula with Frank cluster copulas is very parsimonious and allows for closed-form calculations and very efficient simulations due to its Archimedean clusters.

5 Conclusion

In this paper I introduced the new class of hierarchical Kendall copulas. By grouping variables in clusters, this copula construction method overcomes the curse of dimensionality and thus allows for high-dimensional applications. The use of arbitrary cluster and nesting copulas and potentially several nesting levels allow for very flexible models. Appropriate statistical inference techniques have been presented in detail and illustrated in a substantial application to German stock returns.

Open research questions particularly involve efficient sampling methods for non-Archimedean copulas, since rejection sampling is rather slow. In particular, for the popular class of elliptical copulas, this is an important challenge. Alternative estimation and model selection techniques are also subject of future research.

Acknoweledgement

I like to thank my supervisor Claudia Czado as well as Harry Joe, Matthias Scherer and Michael Smith for helpful remarks and discussions. Furthermore I gratefully acknowledge the support of the TUM Graduate School's International School of Applied Mathematics as well as of Allianz Deutschland AG. The numerical computations were performed on a Linux cluster supported by DFG grant INST 95/919-1 FUGG.

A Proofs

Proof of Theorem 3.5: The idea is to derive the conditional density $f_{j|1,...,j-1}(\cdot|u_1,...,u_{j-1},z)$ and then integrate to obtain the distribution function. One begins by observing that

$$f_{j|1,...,j-1}(u_j|u_1,...,u_{j-1},z) = \frac{f(u_1,...,u_{j-1},u_j,z)}{f(u_1,...,u_{j-1},z)}. \tag{A.1}$$

The numerator can then be rewritten as

$$\begin{aligned}
&f(u_1,...,u_j,z) = P(U_1 = u_1,...,U_j = u_j, C(\boldsymbol{U}) = z)\\
&= \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} P(U_1 = u_1,...,U_{d-1} = u_{d-1}, C(\boldsymbol{U}) = z)\ du_{d-1}...du_{j+1}\\
&= \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} P(U_1 = u_1,...,U_{d-1} = u_{d-1}, U_d = C^{-1}_{u_1,...,u_{d-1}}(z))\ du_{d-1}...du_{j+1}\\
&= \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} c(u_1,...,u_{d-1}, C^{-1}_{u_1,...,u_{d-1}}(z)) \frac{\partial}{\partial z} C^{-1}_{u_1,...,u_{d-1}}(z)\ du_{d-1}...du_{j+1}\\
&= g(u_j),
\end{aligned}$$

where g is defined in (3.2).

Further the denominator of (A.1) then reads as[4]

$$f(u_1, ..., u_{j-1}, z) = \int_{C^{-1}_{u_1,...,u_{j-1}}(z)}^{1} g(u_j)\ du_j.$$

By integration one then obtains the expression for the conditional distribution function given in (3.1). $\square$

Proof of Lemma 3.6: First observe that the density of an exchangeable Archimedean copula C with copula distribution function

$$C(u_1, ..., u_d) = \varphi^{-1}(\varphi(u_1) + ... + \varphi(u_d)) \tag{A.2}$$

only depends on $u_1, ..., u_d$ through the first derivatives of φ and through $C(u_1, ..., u_d)$:

$$c(u_1, ..., u_d) = (\varphi^{-1})^{(d)}(\varphi(u_1) + ... + \varphi(u_d)) \prod_{i=1}^{d} \varphi'(u_i) =: h(C(u_1, ..., u_d)) \prod_{i=1}^{d} \varphi'(u_i). \tag{A.3}$$

To see this observe that any derivative of an inverse is a function of derivatives of the original function applied to the inverse, i.e., $(f^{-1})^{(d)}(x) = \tilde{f}(f^{-1}(x)),\ d \in \mathbb{N}$, for an appropriately chosen $\tilde{f}$. The rest follows from the definition of exchangeable Archimedean copulas (A.2).

Further for exchangeable Archimedean copulas the inverse copula distribution function C^{-1} can easily be determined in closed form as (see Example 2.2)

$$C^{-1}_{u_1,...,u_{d-1}}(z) = \varphi^{-1}\left(\varphi(z) - \sum\nolimits_{1\le i<d} \varphi(u_i)\right), \tag{A.4}$$

and its derivative as

$$\frac{\partial}{\partial z} C^{-1}_{u_1,...,u_{d-1}}(z) = \frac{\varphi'(z)}{\varphi'(\varphi^{-1}(\varphi(z) - \sum_{1\le i<d} \varphi(u_i)))}. \tag{A.5}$$

Following Theorem 3.5 one then determines the function g given in (3.2) by plugging in Expressions (A.3)-(A.5):

$$\begin{aligned} g(u_j) &= \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} c(u_1, ..., u_{d-1}, C^{-1}_{u_1,...,u_{d-1}}(z)) \frac{\partial}{\partial z} C^{-1}_{u_1,...,u_{d-1}}(z)\ du_{d-1}...du_{j+1} \\ &= \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} h(z) \prod_{i=1}^{d-1} \varphi'(u_i) \varphi'(z)\ du_{d-1}...du_{j+1} \\ &= h(z)\varphi'(z) \prod_{i=1}^{j} \varphi'(u_i) \int_{C^{-1}_{u_1,...,u_j}(z)}^{1} \varphi'(u_{j+1}) ... \int_{C^{-1}_{u_1,...,u_{d-2}}(z)}^{1} \varphi'(u_{d-1})\ du_{d-1}...du_{j+1}, \end{aligned} \tag{A.6}$$

since

$$\varphi'(C^{-1}_{u_1,...,u_{d-1}}(z)) \frac{\partial}{\partial z} C^{-1}_{u_1,...,u_{d-1}}(z) = \varphi'(z),$$

and

$$h(C(u_1, ..., u_{d-1}, C^{-1}_{u_1,...,u_{d-1}}(z))) = h(z).$$

---

[4] In the case $j = 1$ this is the density expression of the Kendall distribution function (2.1).

Next, one iteratively solves the nested integrals in (A.6). First,

$$\int_{C^{-1}_{u_1,\dots,u_{d-2}}(z)}^{1} \varphi'(u_{d-1})du_{d-1} = \varphi(1) - \varphi(C^{-1}_{u_1,\dots,u_{d-2}}(z)) \underset{(\text{A.4})}{\overset{\varphi(1)=0}{=}} -\left(\varphi(z) - \sum\nolimits_{1\le i<d-1}\varphi(u_i)\right)$$

$$= \varphi(u_{d-2}) + \left(\sum\nolimits_{1\le i<d-2}\varphi(u_i) - \varphi(z)\right).$$

The second integral is then given by

$$\int_{C^{-1}_{u_1,\dots,u_{d-3}}(z)}^{1} \varphi'(u_{d-2})\left(\varphi(u_{d-2}) + \left(\sum\nolimits_{1\le i<d-2}\varphi(u_i) - \varphi(z)\right)\right)du_{d-2}$$

$$= \left[\frac{1}{2}\left(\varphi(u_{d-2}) + \left(\sum\nolimits_{1\le i<d-2}\varphi(u_i) - \varphi(z)\right)\right)^2\right]_{u_{d-2}=C^{-1}_{u_1,\dots,u_{d-3}}(z)}^{u_{d-2}=1}$$

$$= \frac{1}{2}\left(\varphi(u_{d-3}) + \left(\sum\nolimits_{1\le i<d-3}\varphi(u_i) - \varphi(z)\right)\right)^2.$$

Similarly, the third integral computes to

$$\int_{C^{-1}_{u_1,\dots,u_{d-4}}(z)}^{1} \varphi'(u_{d-3})\frac{1}{2}\left(\varphi(u_{d-3}) + \left(\sum\nolimits_{1\le i<d-3}\varphi(u_i) - \varphi(z)\right)\right)^2 du_{d-3}$$

$$= \dots = \frac{1}{2}\frac{1}{3}\left(\varphi(u_{d-4}) + \left(\sum\nolimits_{1\le i<d-4}\varphi(u_i) - \varphi(z)\right)\right)^3$$

By continuing iteratively, one finally arrives at

$$g(u_j) = h(z)\varphi'(z)\prod_{i=1}^{j}\varphi'(u_i)\frac{1}{(d-j-1)!}\left(\sum\nolimits_{1\le i<j+1}\varphi(u_i) - \varphi(z)\right)^{d-j-1},$$

and thus

$$\int_{C^{-1}_{u_1,\dots,u_{j-1}}(z)}^{u} g(u_j)\ du_j$$

$$= \frac{1}{(d-j-1)!}h(z)\varphi'(z)\prod_{i=1}^{j-1}\varphi'(u_i)\int_{C^{-1}_{u_1,\dots,u_{j-1}}(z)}^{u}\varphi'(u_j)\left(\varphi(u_j) + \left(\sum\nolimits_{1\le i<j}\varphi(u_i) - \varphi(z)\right)\right)^{d-j-1}du_j$$

$$= \frac{1}{(d-j-1)!}h(z)\varphi'(z)\prod_{i=1}^{j-1}\varphi'(u_i)\frac{1}{d-j}\left(\varphi(u) + \sum\nolimits_{1\le i<j}\varphi(u_i) - \varphi(z)\right)^{d-j}$$

$$= \frac{1}{(d-j)!}h(z)\varphi'(z)\prod_{i=1}^{j-1}\varphi'(u_i)\left(\varphi(u) + \sum\nolimits_{1\le i<j}\varphi(u_i) - \varphi(z)\right)^{d-j}. \tag{A.7}$$

By plugging $u = 1$ into (A.7), one further obtains

$$\int_{C^{-1}_{u_1,\dots,u_{j-1}}(z)}^{1} g(u_j)du_j = \frac{1}{(d-j)!}h(z)\varphi'(z)\prod_{i=1}^{j-1}\varphi'(u_i)\left(\sum\nolimits_{1\le i<j}\varphi(u_i) - \varphi(z)\right)^{d-j}. \tag{A.8}$$

Combining Equations (A.7) and (A.8) as in (3.1) then gives

$$F_{j|1,\dots,j-1}(u|u_1,\dots,u_{j-1},z) = \frac{\int_{C^{-1}_{u_1,\dots,u_{j-1}}(z)}^{u} g(u_j)du_j}{\int_{C^{-1}_{u_1,\dots,u_{j-1}}(z)}^{1} g(u_j)du_j} = \frac{\left(\varphi(u) + \sum_{1\le i<j}\varphi(u_i) - \varphi(z)\right)^{d-j}}{\left(\sum_{1\le i<j}\varphi(u_i) - \varphi(z)\right)^{d-j}}$$

$$= \left(1 - \frac{\varphi(u)}{\varphi(z) - \sum_{1\le i<j}\varphi(u_i)}\right)^{d-j},$$

which is the desired result. □

References

Anjos, U. d. and N. Kolev (2005). An application of Kendall distributions. *Review of Business and Economics L*(1), 95–102.

Arbenz, P., C. Hummel, and G. Mainik (2011). Copula based hierarchical risk aggregation through sample reordering. *Forthcoming in Insurance: Mathematics and Economics*.

Barbe, P., C. Genest, K. Ghoudi, and B. Rémillard (1996). On Kendall's process. *Journal of Multivariate Analysis 58*(2), 197–229.

Candelon, B., G. Colletaz, C. Hurlin, and S. Tokpavi (2011). Backtesting Value-at-Risk: A GMM duration-based test. *Journal of Financial Econometrics 9*(2), 314–343.

Chakak, A. and M. Ezzerg (2000). Bivariate contours of copula. *Communications in Statistics - Simulation and Computation 29*(1), 175–185.

Chakak, A. and L. Imlahi (2001). Multivariate probability integral transformation: application to maximum likelihood estimation. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A: Matemáticas 95*(2), 201–212.

Cherubini, U., E. Luciano, and W. Vecchiato (2004). *Copula Methods in Finance*. Chichester: John Wiley & Sons.

Christoffersen, P. F. (1998). Evaluating interval forecasts. *International Economic Review 39*, 841–862.

Christoffersen, P. F. and D. Pelletier (2004). Backtesting Value-at-Risk: A duration-based approach. *Journal of Financial Econometrics 2*(1), 84–108.

Daul, S., E. De Giorgi, F. Lindskog, and A. McNeil (2003). The grouped t-copula with an application to credit risk. *Risk 16*(11), 73–76.

Devroye, L. (1986). *Non-Uniform Random Variate Generation*. New York: Springer.

Genest, C., J. Nešlehová, and J. Ziegel (2011). Inference in multivariate Archimdean copulas. *Test 20*, 223–292.

Genest, C., J. J. Quesada Molina, and J. A. Rodríguez Lallena (1995). De l'impossibilité de construire des lois à marges multidimensionnelles données à partir de copules. *Comptes Rendus de l'Académie des Sciences, Series I, Mathematics 320*, 723–726.

Genest, C. and L.-P. Rivest (1993). Statistical inference procedures for bivariate Archimedean copulas. *Journal of the American Statistical Association 88*(423), 1034–1043.

Genest, C. and L.-P. Rivest (2001). On the multivariate probability integral transformation. *Statistics & Probability Letters 53*(4), 391–399.

Haas, M. (2001). New methods in backtesting. Technical report, Research Center Caesar.

Hastie, T., R. Tibshirani, and J. H. Friedman (2009). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction* (2nd ed.). New York: Springer.

Hering, C. (2011). *Estimation techniques and goodness-of-fit tests for certain copula classes in large dimensions*. Ph. D. thesis, Universität Ulm.

Hofert, M. (2010). Construction and sampling of nested archimedean copulas. In P. Jaworski, F. Durante, W. Härdle, and T. Rychlik (Eds.), *Copula Theory and Its Applications*. Berlin: Springer.

Imlahi, L., M. Ezzerg, and A. Chakak (1999). Estimación de la curva mediana de una cópula $C(x_1, ..., x_m)$. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A: Matemáticas 93*(2), 241–250.

Joe, H. (1997). *Multivariate Models and Dependence Concepts*. London: Chapman & Hall.

Kupiec, P. H. (1995). Techniques for verifying the accuracy of risk measurement models. *The Journal of Derivatives 3*(2), 73–84.

Li, Q. and J. S. Racine (2006). *Nonparametric Econometrics: Theory and Practice*. Princeton: Princeton University Press.

Marco, J. M. and C. Ruiz-Rivas (1992). On the construction of multivariate distributions with given nonoverlapping multivariate marginals. *Statistics & Probability Letters 15*, 259–265.

McNeil, A. J. and J. Nešlehová (2009). Multivariate Archimedean copulas, d-monotone functions and ℓ_1-norm symmetric distributions. *Annals of Statistics 37*(5B), 3059–3097.

Nelsen, R. B. (2006). *An Introduction to Copulas* (2nd ed.). Berlin: Springer.

Nelsen, R. B., J. J. Quesada Molina, J. A. Rodríguez Lallena, and M. Úbeda Flores (2003). Kendall distribution functions. *Statistics & Probability Letters 65*(3), 263–268.

Savu, C. and M. Trede (2010). Hierarchical Archimedean copulas. *Quantitative Finance 10*, 295–304.

Wang, W. and M. T. Wells (2000). Model selection and semiparametric inference for bivariate failure-time data. *Journal of the American Statistical Association 95*, 62–72.

Wu, F., E. Valdez, and M. Sherris (2007). Simulating from exchangeable Archimedean copulas. *Communications in Statistics - Simulation and Computation 36*(5), 1019–1034.

Xue, L. and J. Wang (2010). Distribution function estimation by constrained polynomial spline regression. *Journal of Nonparametric Statistics 22*(4), 443–457.